# ATTACHMENT 2 IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

William SCOTT et al. Confirmation No.: 9834

Serial No.: 09/948,915 Examiner: Rita J. Desai

Filed: September 10, 2001 Group Art Unit: 1625

Title: ω-CARBOXYARYL SUBSTITUTED DIPHENYL UREAS AS RAF KINEASE

**INHIBITORS** 

# **SECOND NON-RESPONSIVE AMENDMENT**

Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

Sir:

To eliminate an obviousness type double patenting rejection in co-pending Application No. 09/899,227, kindly amend the above-identified application to cancel claim 61.

Amendments to the claims begin on page 2 of this paper.

Remarks/Arguments begin on page 50 of this paper.

This listing of claims will replace all prior versions, and listings, of claims in the application:

# **LISTING OF CLAIMS:**

or a pharmaceutically acceptable salt thereof, wherein

D is -NH-C(O)-NH-,

A is a substituted moiety of up to 40 carbon atoms of the formula:  $-L-(M-L^1)_q$ , where L is a 6 membered cyclic structure, which is substituted phenyl or unsubstituted phenyl, bound directly to D,  $L^1$  comprises a substituted cyclic moiety having at least 5 members which is phenyl or pyridinyl,

M is a bridging group which comprises -O-, -S-, or -NR<sup>7</sup>- wherein R<sup>7</sup> is hydrogen, q is an integer of from 1-3; and

B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms with at least one 6-member cyclic structure bound directly to D which is phenyl or pyridinyl,

wherein  $L^1$  is substituted by  $-C(O)R_x$ ,

,  $C_{3-10}$  cycloalkyl having 0-3 heteroatoms,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_3$  -  $C_{12}$  hetaryl having 1-3 heteroatoms selected from, S, N and O,  $C_{7-24}$  alkaryl,  $C_{7-24}$  or aralkyl, and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl up to per halo cycloalkyl having 0-3 heteroatoms selected from N, S and O, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl having 1-3 heteroatoms selected from O, N and S, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C(O)R_g$ ;

 $R_x$  is  $R_z$  or  $NR_aR_b$  where  $R_a$  and  $R_b$  are

a) independently hydrogen,

a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from O, N and S,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O,  $C_{7-24}$  aralkyl, or  $C_7$ - $C_{24}$  alkaryl, and is optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C(O)R_g$  and are optionally substituted by halogen, or

-OSi( $R_f$ )3 where  $R_f$  is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{6-12}$  aryl,  $C_{3}$ - $C_{12}$  hetaryl having 1-3 heteroatoms selected from O, S and N, or  $C_{7-24}$  aralkyl, and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C_{10}$ 0,  $C_{10}$ 0,  $C_{11}$ 0, alkoxyl, halo substituted  $C_7$ - $C_{12}$ 0, alkaryl up to per halo alkaryl, or - $C_{12}$ 0, alkaryl up to per halo alkaryl, or - $C_{12}$ 0, and  $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 1, alkaryl up to per halo alkaryl, or - $C_{12}$ 2, aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{12}$ 3 alkaryl up to per halo alkaryl, or - $C_{12}$ 3, aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{12}$ 4 alkaryl up to per halo alkaryl, or - $C_{12}$ 4 aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{12}$ 4 alkaryl up to per halo alkaryl, or - $C_{12}$ 4 alkaryl up to p

b)  $R_a$  and  $R_b$  together form a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, optionally substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl, halo substituted  $C_{1-6}$  alkyl up to per halo alkyl, halo

substituted  $C_6$ - $C_{12}$  aryl up to per halo aryl, halo substituted  $C_3$ - $C_{12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C(O)R_g$ , or

one of  $R_a$  or  $R_b$  is -C(O)-, a  $C_1$ - $C_5$  divalent alkylene group or a substituted  $C_1$ - $C_5$  divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted  $C_1$ - $C_5$  divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo alkaryl, or - $C(O)R_g$ ,

where B is substituted, L is substituted or  $L^1$  is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and Wn, where n is 0-3;

wherein each W is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -Q-Ar, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, S and N, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>7</sub>-C<sub>24</sub> alkaryl, C<sub>7</sub>-C<sub>24</sub> aralkyl, C<sub>3</sub>-C<sub>12</sub> heteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup> and halogen up to per-halo; with each R<sup>7</sup> independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, S and N, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>3</sub>-C<sub>13</sub> hetaryl having 1-3

heteroatoms selected from O, N and S, C<sub>7</sub>-C<sub>14</sub> alkaryl, C<sub>7</sub>-C<sub>24</sub> aralkyl, or C<sub>4</sub>-C<sub>23</sub> alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by halogen,

wherein Q is -O-, -S-, -N(R<sup>7</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>7</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>- CHX<sup>a</sup>-, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>7</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m= 1-3, and  $X^a$  is halogen; and

Ar is a 5- or 6-member aromatic structure containing 0-2 members selected from the group consisting of nitrogen, oxygen and sulfur, which is optionally substituted by halogen, up to per-halo, and optionally substituted by  $Z_{n1}$ , wherein n1 is 0 to 3 and each Z is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup> -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which is C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6</sub>-C<sub>14</sub> aryl, or C<sub>3</sub>-C<sub>13</sub> hetaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and -NR<sup>7</sup>C(O)OR<sup>7</sup>, with R<sup>7</sup> as defined above

where  $R_g$  is  $C_{1-10}$  alkyl; -CN, -CO<sub>2</sub>R<sub>d</sub>, -OR<sub>d</sub>, -SR<sub>d</sub>, -NO<sub>2</sub>, -C(O)R<sub>e</sub>, -NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub> C(O)OR<sub>e</sub> and -NR<sub>d</sub> C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen,  $C_{1-10}$ , alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl having 0-3 heteroatoms selected from O, N and S,  $C_{6-12}$  aryl,  $C_3$ -C<sub>12</sub> hetaryl with 1-3 heteroatoms selected from O, N and S and  $C_7$ -C<sub>24</sub> aralkyl,  $C_7$ -C<sub>24</sub> alkaryl, up to per halo substituted  $C_1$ -C<sub>10</sub> alkyl, up to per halo substituted  $C_3$ -C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per halo substituted  $C_6$ -C<sub>14</sub> aryl, up to per halo substituted  $C_3$ -C<sub>12</sub> hetaryl having 1-3 heteroatoms selected from O, N, and S, halo substituted  $C_7$ -C<sub>24</sub> alkaryl up to per halo alkaryl, and up to per halo substituted  $C_7$ -C<sub>24</sub> aralkyl.

# **Claim 2.** (**Original**) A compound as in claim 1 wherein:

 $R_y$  is hydrogen,  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl having 0-3 heteroatoms,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{7-24}$  aralkyl,  $C_{7-24}$  alkaryl, substituted  $C_{1-10}$  alkyl, substituted  $C_{1-10}$  alkoxy, substituted  $C_{3-10}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O, substituted  $C_{6}$ - $C_{14}$  aryl, substituted  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,

substituted  $C_{7-24}$  alkaryl or substituted  $C_7-C_{24}$  aralkyl, where  $R_y$  is a substituted group, it is substituted by halogen up to per halo,

 $R_z$  is hydrogen,  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl having 0-3 heteroatom,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_3$  - $C_{12}$  hetaryl having 1-3 heteroatoms selected from, S, N and O,  $C_{7-24}$  alkaryl,  $C_{7-24}$  aralkyl, substituted  $C_{1-10}$  alkyl, substituted  $C_{1-10}$  alkoxy, substituted  $C_{6}$ - $C_{14}$  aryl, substituted  $C_{3}$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from S, N and O, substituted  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from S, N and O, substituted  $C_{7-24}$  aralkyl where  $R_z$  is a substituted group, it is substituted by halogen up to per halo, hydroxy,  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_{6}$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_{3}$ - $C_{12}$  halo substituted  $C_{3}$ - $C_{12}$  hetaryl up to per halo hetaryl having 1-3 heteroatoms selected from O, N and S, halo substituted  $C_{7}$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_{7}$ - $C_{24}$  alkaryl up to per halo alkaryl, and - $C(O)R_g$ ,

R<sub>a</sub> and R<sub>b</sub> are,

# a) independently hydrogen,

a carbon based moiety selected from the group consisting of  $C_1$  - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from O, N and S,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O,  $C_{7-24}$  aralkyl,  $C_7$ - $C_{24}$  alkaryl, substituted  $C_{1-10}$  alkyl, substituted  $C_{1-10}$  alkoxy, substituted  $C_{3-10}$  cycloalkyl, having 0-3 heteroatoms selected from N, S and O, substituted  $C_{6-12}$  aryl, substituted  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O, substituted  $C_{7-24}$  aralkyl, substituted  $C_{7-24}$  alkaryl, where  $R_a$  and  $R_b$  are a substituted group, they are substituted by halogen up to per halo, hydroxy,  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo heteraryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, and - $C(O)R_g$ ; or

or

b)  $R_a$  and  $R_b$  together form a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, or a substituted 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O with substituents selected from the group consisting of halogen up to per halo, hydroxy,  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl, halo substituted  $C_{1-6}$  alkyl up to per halo alkyl, halo substituted  $C_6$ - $C_{12}$  aryl up to per halo aryl, halo substituted  $C_3$ - $C_{12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo heteraryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, and - $C(O)R_g$ ,

or

c) one of  $R_a$  or  $R_b$  is -C(O)-, a  $C_1$ - $C_5$  divalent alkylene group or a substituted  $C_1$ - $C_5$  divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members,

wherein the substituents of the substituted  $C_1$ - $C_5$  divalent alkylene group are selected from the group consisting of halogen, hydroxy,  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$ - $C_{24}$  alkaryl,  $C_7$ - $C_{24}$  aralkyl,  $C_{1-6}$  halo substituted alkyl

up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo heteraryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, and - $C(O)R_g$ ,

where  $R_g$  is  $C_{1-10}$  alkyl; -CN, -CO<sub>2</sub>R<sub>d</sub>, -OR<sub>d</sub>, -SR<sub>d</sub>, -NO<sub>2</sub>, -C(O)R<sub>e</sub>, -NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub> C(O)OR<sub>e</sub> and -NR<sub>d</sub> C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen,  $C_{1-10}$ , alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl having 0-3 heteroatoms selected from O, N and S,  $C_{6-12}$  aryl,  $C_3$ -C<sub>12</sub> hetaryl with 1-3 heteroatoms selected from O, N and S and  $C_7$ -C<sub>24</sub> aralkyl,  $C_7$ -C<sub>24</sub> alkaryl, up to per halo substituted  $C_1$ -C<sub>10</sub> alkyl, up to per halo substituted  $C_3$ -C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per halo substituted  $C_6$ -C<sub>14</sub> aryl, up to per halo substituted  $C_3$ -C<sub>12</sub> hetaryl having 1-3 heteroatoms selected from O, N, and S, halo substituted  $C_7$ -C<sub>24</sub> alkaryl up to per halo alkaryl, and up to per halo substituted  $C_7$ -C<sub>24</sub> aralkyl,

W is independently selected from the group consisting of -CN, -CO $_2$ R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO $_2$ , -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, S and N, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>7</sub>-C<sub>24</sub> alkaryl, C<sub>7</sub> -C<sub>24</sub> aralkyl, C<sub>3</sub>-C<sub>12</sub> heteroaryl having 1-3 heteroatoms selected from O, N and S, C<sub>4</sub>-C<sub>23</sub> alkheteroaryl having 1-3 heteroatoms selected from O, N and S, substituted C<sub>1</sub>-C<sub>10</sub> alkyl, substituted C<sub>1</sub>-C<sub>10</sub> alkoxy, substituted C<sub>2</sub>-C<sub>10</sub> alkenyl, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, substituted C<sub>6</sub>-C<sub>12</sub> aryl, substituted C<sub>3</sub>-C<sub>12</sub> hetaryl having 1-3 heteroatoms selected from O, N and S, substituted C<sub>7</sub>-C<sub>24</sub> aralkyl, substituted C<sub>7</sub>-C<sub>24</sub> alkaryl, substituted C<sub>4</sub>-C<sub>23</sub> alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and -Q-Ar;

 $R^7$  is independently selected from H,  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenoyl,  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_6$ - $C_{14}$  aryl,  $C_3$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S,  $C_7$ - $C_{14}$  alkaryl,  $C_7$ - $C_{24}$  aralkyl,  $C_4$ - $C_{23}$  alkheteroaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_1$ - $C_{10}$  alkyl, up to perhalosubstituted  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_6$ - $C_{14}$  aryl, up to perhalosubstituted  $C_7$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_{14}$  aryl, up to perhalosubstituted  $C_7$ - $C_{15}$  hetaryl having 1-3 heteroatoms selected from O, N and S, up to perhalosubstituted  $C_7$ - $C_$ 

halosubstituted  $C_7$ - $C_{24}$  aralkyl, up to per-halosubstituted  $C_7$ - $C_{24}$  alkaryl, and up to per-halosubstituted  $C_4$ - $C_{23}$  alkheteroaryl; and

each Z is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, - SR<sup>7</sup> -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>3</sub>-C<sub>13</sub> hetaryl having 1-3 heteroatoms selected from O, N and S, substituted C<sub>1</sub>-C<sub>24</sub> aralkyl, C<sub>4</sub>-C<sub>23</sub> alkheteroaryl having 1-3 heteroatoms selected from O, N and S, substituted C<sub>1</sub>-C<sub>10</sub> alkyl, substituted C<sub>1</sub>-C<sub>10</sub> alkoxy, substituted C<sub>2</sub>-C<sub>10</sub> alkenyl, substituted C<sub>1</sub>-C<sub>10</sub> alkenoyl, substituted C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, substituted C<sub>6</sub>-C<sub>12</sub> aryl, substituted C<sub>7</sub>-C<sub>24</sub> alkaryl, substituted C<sub>7</sub>-C<sub>24</sub> aralkyl and substituted C<sub>4</sub>-C<sub>23</sub> alkheteroaryl having 1-3 heteroatoms selected from O, N and S; wherein if Z is a substituted group, the one or more substituents are selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -CO()NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and -NR<sup>7</sup>C(O)OR<sup>7</sup>.

Claim 3. (Previously Presented) A compound as in claim 1 wherein M is -O-.

**Claim 4.** (**Previously Presented**) A compound as in claim 1 wherein the cyclic structures of B and L bound directly to D have hydrogen substituents in the ortho position.

Claim 5. (Canceled)

Claim 6. (Canceled)

Claim 7. (Previously Presented) A compound of claim 1 wherein B of Formula I is an unsubstituted phenyl group.

Claim 8. (Previously Presented) A compound of claim 1 wherein B of Formula I is a substituted phenyl group, substituted 1 to 3 times by 1 or more substituents

selected from the group consisting of -CN, halogen,  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy, -OH, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkoxy or phenyl substituted by halogen up to per halo.

# Claim 9. (Canceled)

**Claim 10.** (**Previously presented**) A compound of claim 8, wherein L, the 6 member cyclic structure bound directly to D, is an unsubstituted phenyl group.

# Claim 11. (Canceled)

Claim 12. (Previously Presented) A compound of claim 1, wherein said substituted cyclic moiety  $L^1$  is pyridinyl.

Claim 13. (Previously Presented) A compound of claim 3, wherein said substituted cyclic moiety  $L^1$  is pyridinyl.

Claim 14. (Previously Presented) A compound of claim 7, wherein said substituted cyclic moiety  $L^1$  is pyridinyl.

Claim 15. (Previously Presented) A compound of claim 8, wherein said substituted cyclic moiety  $L^1$  is pyridinyl.

# Claim 16. (Cancelled)

Claim 17. (Previously Presented) A compound of claim 10, wherein said substituted cyclic moiety  $L^1$  is pyridinyl.

Claim 18. (Previously Presented) A compound of claim 14, wherein M is - O-.

Claim 19. (Previously Presented) A compound of claim 15, wherein M is - O-.

# Claim 20. (Cancelled)

**Claim 21.** (**Previously Presented**) A compound of claim 17, wherein M is -O-.

Claim 22. (Original) A compound of claim 1 wherein  $L^1$  is additionally substituted 1 to 3 times by one or more substituents selected from the group consisting of  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, -CN, -OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

Claim 23. (Original) A compound of claim 13 wherein  $L^1$  is additionally substituted 1 to 3 times by one or more substituents selected from the group consisting of  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, -CN, -OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

Claim 24. (Original) A compound of claim 18 wherein  $L^1$  is additionally substituted 1 to 3 times by one or more substituents selected from the group consisting of  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, -CN, -OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

Claim 25. (Original) A compound of claim 19 wherein  $L^1$  is additionally substituted 1 to 3 times by one or more substituents selected from the group consisting of  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, -CN, -OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

# Claim 26. (Canceled)

Claim 27. (Original) A compound of claim 21 wherein  $L^1$  is additionally substituted 1 to 3 times by one or more substituents selected from the group consisting of  $C_1$ -

 $C_{10}$  alkyl, up to per halo substituted  $C_1$ - $C_{10}$  alkyl, -CN, -OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

Claim 28. (Cancelled)

Claim 29. (Cancelled)

Claim 30. (Original) A compound of claim 1 wherein  $L^1$  is substituted only by  $-C(O)R_x$ .

Claim 31. (Cancelled)

Claim 32. (Cancelled)

Claim 33. (Previously Presented) A compound of claim 13 wherein  $L^1$  is substituted by  $-C(O)R_x$  wherein  $R_x$  is  $NR_aR_b$ .

Claim 34. (Previously Presented) A compound of claim 18 wherein  $L^1$  is substituted by  $-C(O)R_x$  wherein  $R_x$  is  $NR_aR_b$ .

Claim 35. (Previously Presented) A compound of claim 19 wherein  $L^1$  is substituted by  $-C(O)R_x$ , wherein  $R_x$  is  $NR_aR_b$ .

Claim 36. (Cancelled)

Claim 37. (Previously Presented) A compound of claim 21 wherein  $L^1$  is substituted by  $-C(O)R_x$  wherein  $R_x$  is  $NR_aR_b$ .

**Claim 38.** (**Previously Presented**) A compound of Formula I:

 $A - D - B \tag{I}$ 

or a pharmaceutically acceptable salt thereof, wherein

D is -NH-C(O)-NH-,

A is a substituted moiety of up to 40 carbon atoms of the formula:  $-L-(M-L^1)_q$ , where L is a 6 membered aryl moiety which is unsubstituted phenyl bound directly to D, L<sup>1</sup> comprises a substituted cyclic moiety having at least 5 members which is phenyl or pyridinyl, M is -O- and

B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms with at least one 6-member cyclic structure bound directly to D which is phenyl or pyridinyl

wherein  $L^1$  is substituted by  $-C(O)R_x$ 

 $R_x$  is  $NR_aR_b$  where  $R_a$  and  $R_b$  are

a) independently hydrogen,

a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is of  $C_1$  - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_{3-12}$  heteryl having 1-3 heteroatoms selected from O, N and S,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O,  $C_{7-24}$  aralkyl or  $C_7$ - $C_{24}$  alkaryl, and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per hal

-OSi( $R_f$ )<sub>3</sub> where  $R_f$  is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and

O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C(O)R_g$ , or

- b)  $R_a$  and  $R_b$  together form a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, or a substituted 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl, halo substituted  $C_{1-6}$  alkyl up to per halo alkyl, halo substituted  $C_6$ - $C_{12}$  aryl up to per halo aryl, halo substituted  $C_3$ - $C_{12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or - $C(O)R_g$ , or
- one of  $R_a$  or  $R_b$  is -C(O)-, a  $C_1$ - $C_5$  divalent alkylene group or a substituted  $C_1$ - $C_5$  divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted  $C_1$ - $C_5$  divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, halo substituted by halogen;

where B is substituted, L is substituted or  $L^1$  is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and Wn, where n is 0-3;

wherein each W is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -Q-Ar, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected

from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenoyl,  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_6$ - $C_{14}$  aryl,  $C_7$ - $C_{24}$  alkaryl,  $C_7$ - $C_{24}$  aralkyl, or  $C_3$ - $C_{12}$  heteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents independently selected from the group consisting of -CN, - $CO_2R^7$ , - $C(O)R^7$ , - $C(O)NR^7R^7$ , - $OR^7$ , - $SR^7$ , - $NR^7R^7$ , - $NO_2$ , - $NR^7C(O)R^7$ , - $NR^7C(O)OR^7$  and halogen up to per-halo; with each  $R^7$  independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenoyl,  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_6$ - $C_{14}$  aryl,  $C_3$ - $C_{13}$  heteryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by halogen,

wherein Q is -O-, -S-, -N(R<sup>7</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>7</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>- CHX<sup>a</sup>-, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>7</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m= 1-3, and  $X^a$  is halogen;

Ar is a 5- or 6-member aromatic structure containing 0-2 members selected from the group consisting of nitrogen, oxygen and sulfur, which is optionally substituted by halogen, up to per-halo, and optionally substituted by  $Z_{n1}$ , wherein n1 is 0 to 3 and each Z is independently selected from the group consisting of -CN, -CO $_2$ R $^7$ , -C(O)R $^7$ , -C(O)NR $^7$ R $^7$ , -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup> -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which is C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>3</sub>-C<sub>13</sub> hetaryl having 1-3 heteroatoms selected from O, N and S, C7-C24 alkaryl, C7-C24 aralkyl, C4-C23 alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents are selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and -NR<sup>7</sup>C(O)OR<sup>7</sup>, with R<sup>7</sup> as defined above; and where R<sub>g</sub> is C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen, C<sub>1</sub>-<sub>10</sub>, alkyl, C<sub>1-10</sub> alkoxy, C<sub>3-10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6-</sub> 12 aryl, C<sub>3</sub>-C<sub>12</sub> hetaryl with 1-3 heteroatoms selected from O, N and S and C<sub>7</sub>-C<sub>24</sub> aralkyl, C<sub>7</sub>-C<sub>24</sub> alkaryl, up to per halo substituted C<sub>1</sub>-C<sub>10</sub> alkyl, up to per halo substituted C<sub>3</sub> -C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per halo substituted  $C_6$  -  $C_{14}$  aryl, up to per halo substituted  $C_3$  - $C_{12}$  hetaryl having 1-3 heteroatoms selected from O, N, and S, halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, or up to per halo substituted  $C_7$ - $C_{24}$  aralkyl.

# Claim 39. (Previously Presented) A compound of Formula I:

$$A - D - B \tag{I}$$

or a pharmaceutically acceptable salt thereof, wherein

D is -NH-C(O)-NH-,

A is a substituted moiety of up to 40 carbon atoms of the formula:  $-L-(M-L^1)_q$ , where L is a substituted or unsubstituted phenyl or pyridinyl moiety bound directly to D,  $L^1$  comprises a substituted phenyl, or pyridinyl moiety, M is -O- and

B is a substituted or unsubstituted phenyl group bound directly to D, wherein  $L^1$  is substituted by-C(O)R<sub>x</sub>,

 $R_x$  is  $NR_aR_b$  where  $R_a$  and  $R_b$  are

a) independently hydrogen,

a moiety, which is  $C_1$  - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from O, N and S,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from N, S and O,  $C_{7-24}$  aralkyl, or  $C_7$ - $C_{24}$  alkaryl, and optionally substituted by halogen, hydroxy and carbon based substituents which are- $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_{3-12}$  hetaryl having 1-3 heteroatoms selected from N, S and O,  $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted  $C_3$ - $C_{12}$  hetaryl up to per halo hetaryl, halo substituted  $C_7$ - $C_{24}$  aralkyl up to per halo aralkyl, or halo substituted  $C_7$ - $C_{24}$  alkaryl up to per halo alkaryl, and - $C(O)R_g$ ,

where B is substituted, L is substituted or  $L^1$  is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and Wn, where n is 0-3;

wherein each W is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -Q-Ar, and moieties which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenyl,  $C_3$ - $C_{10}$ 

cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_6$ - $C_{14}$  aryl,  $C_7$ - $C_{24}$  alkaryl,  $C_7$ - $C_{24}$  aralkyl,  $C_3$ - $C_{12}$  heteroaryl having 1-3 heteroatoms selected from O, N and S, or  $C_4$ - $C_{23}$  alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents independently selected from the group consisting of -CN, - $CO_2R^7$ , - $C(O)R^7$ , - $C(O)NR^7R^7$ , - $OR^7$ , - $SR^7$ , - $NR^7R^7$ , - $NO_2$ , - $NR^7C(O)R^7$ , - $NR^7C(O)OR^7$  and halogen up to per-halo; with each  $R^7$  independently selected from H or a moiety which is  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenoyl,  $C_3$ - $C_{10}$  cycloalkyl having 0-3 heteroatoms selected from O, S and N,  $C_6$ - $C_{14}$  aryl,  $C_3$ - $C_{13}$  hetaryl having 1-3 heteroatoms selected from O, N and S,  $C_7$ - $C_{14}$  alkaryl,  $C_7$ - $C_{24}$  aralkyl, or  $C_4$ - $C_{23}$  alkheteroaryl having 1-3 heteroatoms selected from O, N and S,

wherein Q is -O-, -S-, -N(R<sup>7</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>7</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>- CHX<sup>a</sup>-, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>7</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m= 1-3, and X<sup>a</sup> is halogen;

Ar is phenyl or pyridinyl which is optionally substituted by halogen, up to per-halo, and optionally substituted by  $Z_{n1}$ , wherein n1 is 0 to 3 and each Z is independently selected from the group consisting of -CN, -CO $_2$ R $^7$ , -C(O)R $^7$ , -C(O)NR $^7$ R $^7$ , -NO $_2$ , -OR $^7$ , - SR $^7$  - $NR^7R^7$ ,  $-NR^7C(O)OR^7$ ,  $-NR^7C(O)R^7$ , and a moiety which is  $C_1-C_{10}$  alkyl,  $C_1-C_{10}$  alkoxy,  $C_2-C_{10}$ C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6</sub>-C<sub>14</sub> aryl, C<sub>3</sub>-C<sub>13</sub> hetaryl having 1-3 heteroatoms selected from O, N and S, C<sub>7</sub>-C<sub>24</sub> alkaryl, C7-C24 aralkyl, or C4-C23 alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and optionally substituted by one or more substituents selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and  $-NR^7C(O)OR^7; \text{ and where } R_g \text{ is } C_{1\text{-}10} \text{ alkyl}; \text{-}CN, \text{-}CO_2R_d, \text{-}OR_d, \text{-}SR_d, \text{-}NO_2, \text{-}C(O)R_e, \text{-}CO_2R_d, \text{-}OR_d, \text{-}OR_d$ NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub> C(O)OR<sub>e</sub> and -NR<sub>d</sub> C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen, C<sub>1-10</sub>, alkyl, C<sub>1-10</sub> alkoxy, C<sub>3-10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, C<sub>6-12</sub> aryl, C<sub>3</sub>-C<sub>12</sub> hetaryl with 1-3 heteroatoms selected from O, N and S and C7-C24 aralkyl, C7 -C24 alkaryl, up to per halo substituted C1-C<sub>10</sub> alkyl, up to per halo substituted C<sub>3</sub> -C<sub>10</sub> cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per halo substituted  $C_6$  - $C_{14}$  aryl, up to per halo substituted  $C_3$  - $C_{12}$  hetaryl having 1-3 heteroatoms selected from O, N, and S, halo substituted C7-C24 alkaryl up to per halo alkaryl, or up to per halo substituted  $C_7$ - $C_{24}$  aralkyl.

**Claim 40.** (**Previously Presented**) A compound as in claim 38 wherein the cyclic structures of B and L bound directly to D have hydrogen substituents in the ortho position.

#### Claim 41. (Cancelled)

**Claim 42.** (**Previously Presented**) A compound as in claim 39 wherein the cyclic structures of B and L bound directly to D have hydrogen substituents in the ortho position.

#### Claim 43. (Cancelled)

Claim 44. (Original) A compound as in claim 38 wherein substituents for B and L and additional substituents for  $L^1$ , are selected from the group consisting of  $C_1$ - $C_{10}$  alkyl up to per halo substituted  $C_1$ - $C_{10}$  alkyl, CN, OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

Claim 45. (Original) A compound as in claim 39 wherein substituents for B and L and additional substituents for  $L^1$ , are selected from the group consisting of  $C_1$ - $C_{10}$  alkyl up to per halo substituted  $C_1$ - $C_{10}$  alkyl, CN, OH, halogen,  $C_1$ - $C_{10}$  alkoxy and up to per halo substituted  $C_1$ - $C_{10}$  alkoxy.

#### Claim 46. (Cancelled)

#### Claim 47. (Cancelled)

Claim 48. (Previously Presented) A compound of claim 38 wherein  $R_a$  and  $R_b$  are independently hydrogen and  $C_{1-6}$  alkyl.

Claim 49. (Previously Presented) A compound of claim 39 wherein and  $R_a$  and  $R_b$  are independently hydrogen and  $C_{1-6}$  alkyl, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen.

- Claim 50. (Previously Presented) A pharmaceutically acceptable salt of a compound of formula I of claim 1 which is
- a) a basic salt of an organic acid or inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or
- b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.
- Claim 51. (Previously Presented) A pharmaceutically acceptable salt of a compound of claim 61 which is
- a) a basic salt of an organic acid or inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or
- b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.
- Claim 52. (Previously Presented) A pharmaceutically acceptable salt of a compound of claim 33 which is
- a) a basic salt of an organic acid or inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or

b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.

# Claim 53. (Previously Presented) A pharmaceutically acceptable salt of a compound of claim 38 which is

- a) a basic salt of an organic acid or inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or
- b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.

# Claim 54. (Previously Presented) A pharmaceutically acceptable salt of a compound of claim 39 which is

- a) a basic salt of an organic acid or inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or
- b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.
- Claim 55. (Original) A pharmaceutical composition comprising a compound of claim 1 or a pharmaceutically acceptable salt of a compound of formula I, and a physiologically acceptable carrier.

**Claim 56.** (**Previously Presented**) A pharmaceutical composition comprising a compound of claim 2 and a physiologically acceptable carrier.

**Claim 57.** (**Previously Presented**) A pharmaceutical composition comprising a compound of claim 33 and a physiologically acceptable carrier.

**Claim 58.** (**Previously Presented**) A pharmaceutical composition comprising a compound of claim 38 and a physiologically acceptable carrier.

**Claim 59.** (**Previously Presented**) A pharmaceutical composition comprising a compound of claim 39 and a physiologically acceptable carrier.

Claim 60. (Cancelled)

Claim 61. (Cancelled)

**Claim 62.** (**Previously Presented**) A method for the treatment of a cancerous cell growth mediated by raf kinase, comprising administering a compound of claim 1.

**Claim 63.** (**Previously Presented**) A method for the treatment of a cancerous cell growth mediated by raf kinase, comprising administering a compound of claim 33.

**Claim 64.** (**Previously Presented**) A method for the treatment of a cancerous cell growth mediated by raf kinase, comprising administering a compound of claim 38.

**Claim 65.** (**Previously Presented**) A method for the treatment of a cancerous cell growth mediated by raf kinase, comprising administering a compound of claim 39.

Claim 66. (Cancelled)

Claim 67. (Cancelled)

# **Claim 68.** (**Previously Presented** ) A compound of Formula I:

$$A - D - B \tag{I}$$

or a pharmaceutically acceptable salt thereof, wherein

D is -NH-C(O)-NH-,

A is a substituted moiety of the formula:

 $-L-M-L^1$ ,

wherein L is

phenyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_5$  linear or branched haloalkyl up to perhalo,  $C_1$ - $C_3$  alkoxy,  $C_1$ - $C_3$  haloalkoxy up to per haloalkoxy, hydroxy, amino,  $C_1$ - $C_3$  alkylamino,  $C_1$ - $C_6$  dialkylamino, halogen, cyano, and nitro;

- L<sup>1</sup> comprises a substituted cyclic moiety which is
- (i) phenyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $R^7$ ,  $OR^7$ ,  $NR^7R^{7'}$ ,  $C(O)R^7$ ,  $C(O)OR^7$ ,  $C(O)NR^7R^{7'}$ ,  $NR^7C(O)R^{7'}$ ,  $NR^7C(O)OR^{7'}$ , halogen, cyano and nitro;
- (ii) pyridinyl optionally substituted with 1-3 substituents independently selected from the group consisting of  $R^7$ ,  $OR^7$ ,  $NR^7R^{7'}$ ,  $C(O)R^7$ ,  $C(O)OR^7$ ,  $C(O)NR^7R^{7'}$ ,  $NR^7C(O)R^{7'}$ ,  $NR^7C(O)OR^{7'}$ , halogen, cyano and nitro;  $C(O)OR^7$ ,  $C(O)NR^7R^{7'}$ ,  $NR^7C(O)R^{7'}$ ,  $NR^7C(O)OR^{7'}$ , halogen, cyano and nitro; and

wherein  $L^1$  is substituted by  $-C(O)R_x$ wherein  $R_x$  is  $R_z$  or  $NR_aR_b$  and  $R_a$  and  $R_b$  are, independently,  $R_z$ 

M is -O-

B is

- (i) phenyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $R^7$ ,  $OR^7$ ,  $NR^7R^{7'}$ ,  $C(O)R^7$ ,  $C(O)OR^7$ ,  $C(O)NR^7R^{7'}$ ,  $NR^7C(O)R^{7'}$ ,  $NR^7C(O)OR^{7'}$  halogen, cyano, and nitro; or
- (ii) pyridinyl optionally substituted with 1-3 substituents independently selected from the group consisting of  $R^7$ ,  $OR^7$ ,  $NR^7R^{7'}$ ,  $C(O)R^7$ ,  $C(O)OR^7$ ,  $C(O)NR^7R^{7'}$ ,  $NR^7C(O)R^{7'}$ ,  $NR^7C(O)OR^{7'}$ , halogen, cyano, and nitro; and

each R7, R7, Rz and Rf is independently

(a) hydrogen,

- (b)  $C_1$ - $C_6$  linear, branched, or cyclic alkyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy and hydroxy;
- (c)  $C_1$ - $C_6$  alkoxy, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy, hydroxy and halogen;
- (d) phenyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy, hydroxy and halogen,
- (e) 5-6 membered monocyclic heteroaryl having 1-4 heteroatoms selected from the group consisting of O, N and S or 8-10 membered bicyclic heteroaryl having 1-6 heteroatoms selected from the group consisting of O, N and S, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy, hydroxy and halogen,
- (f)  $C_1$ - $C_3$  alkyl-phenyl, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy, hydroxy and halogen; and
- (g) up to per–halo substituted  $C_1$ - $C_5$  linear, branched or cyclic alkyl, and where not per-halo substituted, optionally substituted with 1-3 substituents independently selected from the group consisting of  $C_1$ - $C_5$  linear or branched alkyl, up to perhalo substituted  $C_1$ - $C_5$  linear or branched alkyl,  $C_1$ - $C_3$  alkoxy and hydroxy.

Claim 69. (Cancelled)

Claim 70. (Cancelled)

Claim 71. (Cancelled)

Claim 72. (Cancelled)

Claim 73. (Previously Presented) A compound of claim 68 wherein the substituents of the substituted structures of L are selected from the group consisting of methyl, triflouromethyl, ethyl, n-propyl, n-butyl, n-pentyl, i-propyl, t-butyl, methoxy, ethoxy, propoxy, Cl, Br, F, cyano, nitro, hydroxy, amino, methylamino, dimethylamino, ethylamino and diethylamino.

Claim 74. (Previously Presented) A compound of claim 68 wherein the substituents of the substituted structures of B and L<sup>1</sup> are independently selected from the group consisting of methyl, triflouromethyl, ethyl, n-propyl, n-butyl, n-pentyl, isopropyl, tert-butyl, sec-butyl, isobutyl, cyclopropyl, cyclobutyl, cyclopentyl, methoxy, ethoxy, propoxy, Cl, Br and F, cyano, nitro, hydroxy, amino, methylamino, dimethylamino, ethylamino and diethylamino.

#### Claim 75. (Cancelled)

Claim 76. (Previously Presented) A compound as in claim 68 wherein B, L and  $L^1$  follow one of the following of combinations:

B= phenyl, L=phenyl and  $L^1$  is phenyl,

B= phenyl, L=phenyl and L<sup>1</sup> is pyridinyl,

B=pyridinyl, L= phenyl and  $L^1$  is phenyl,

B=pyridinyl, L=phenyl and L1 is pyridinyl,

Claim 77. (Previously Presented) A pharmaceutical composition for the treatment of a cancerous cell growth comprising a compound of claim 68 or a pharmaceutically acceptable salt of a compound of formula I and a physiologically acceptable carrier.

#### Claim 78. (Cancelled)

# Claim 79. (Cancelled)

Claim 80. (Previously Presented) A pharmaceutical composition for the treatment of a cancerous cell growth as in claim 77 wherein the pharmaceutically acceptable salt is

a) a basic salt of an organic acid or an inorganic acid which is hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, trifluoromethanesulfonic acid, benzenesulfonic acid, p-toluene sulfonic acid (tosylate salt), 1-napthalene sulfonic acid, 2-napthalene sulfonic acid, acetic acid, trifluoroacetic acid, malic

acid, tartaric acid, citric acid, lactic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, benzoic acid, salicylic acid, phenylacetic acid, or mandelic acid; or

b) an acid salt of an organic or inorganic base containing an alkali metal cation, an alkaline earth metal cation, an ammonium cation, an aliphatic substituted ammonium cation or an aromatic substituted ammonium cation.

Claim 81. (Cancelled)

Claim 82. (Cancelled)

**Claim 83.** (**Previously presented**) A compound of Formula I:

$$A - D - B \tag{I}$$

or a pharmaceutically acceptable salt thereof, wherein

D is -NH-C(O)-NH-,

A is a substituted moiety of up to 40 carbon atoms of the formula:  $-L-(M-L^1)_q$ , where L is a 6 membered cyclic structure, which is substituted phenyl or unsubstituted phenyl, bound directly to D,  $L^1$  comprises a substituted cyclic moiety having at least 5 members which is phenyl or pyridinyl,

M is one or more bridging groups selected from the group consisting of -O-, -S-, -NR<sup>7</sup>- where R<sup>7</sup> is hydrogen, q is an integer of from 1-3; and

B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms with at least one 6-member cyclic structure bound directly to D which is phenyl or pyridinyl,

wherein  $L^1$  is substituted by  $-S(O)_2R_x$ ,

 $R_z$  is hydrogen or a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenyl,  $C_{6-12}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- oxadiazol-3- o

thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

S, N and O, and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl up to per halo cycloalkyl halo substituted, up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl,

1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or  $-C(O)R_g$ ;

 $R_x$  is  $R_z$  or  $NR_aR_b$  where  $R_a$  and  $R_b$  are

a) independently hydrogen,

a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is C<sub>1</sub> -C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>3-10</sub> cycloalkyl, C<sub>2-10</sub> alkenyl,  $C_{1-10}$  alkenyl,  $C_{6-12}$  aryl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

 $C_{3-12}$  cycloalkyl and is optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5-, or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl, up to per halo cycloalkyl, halo substituted up to per halo 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7-, or 8-quinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7

quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or -C(O)R<sub>g</sub> and are optionally substituted by halogen, or

-OSi(R<sub>f</sub>)<sub>3</sub> where R<sub>f</sub> is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O, which is C<sub>1-10</sub> alkyl, C<sub>1-10</sub> alkoxy, C<sub>3</sub>-C<sub>10</sub> cycloalkyl, C<sub>6-12</sub> aryl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 3-, 4-, 5-, 6- or 7-benzotyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-

, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl, C<sub>3</sub>-C<sub>12</sub> halo substituted cycloalkyl up to per halo cycloalkyl, halo substituted up to per halo 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

or  $-C(O)R_g$ , or

b) R<sub>a</sub> and R<sub>b</sub> together form 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-

benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

optionally substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are C<sub>1</sub>-<sub>10</sub> alkyl, C<sub>3-12</sub> cycloalkyl 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl, halo substituted  $C_{1-6}$  alkyl up to per halo alkyl, halo substituted  $C_6$ - $C_{12}$  aryl up to per halo aryl, halo substituted  $C_3$ - $C_{12}$  cycloalkyl up to per halo cycloalkyl, halo substituted up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-,

3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or  $-C(O)R_g$ , or

one of R<sub>a</sub> or R<sub>b</sub> is -C(O)-, a C<sub>1</sub>-C<sub>5</sub> divalent alkylene group or a substituted C<sub>1</sub>-C<sub>5</sub> divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted C<sub>1</sub>-C<sub>5</sub> divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are C<sub>1-10</sub> alkyl, C<sub>3-12</sub> cycloalkyl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl up to per halo cycloalkyl, halo substituted up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4-

or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 3-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or  $-C(O)R_g$ ,

where B is substituted, L is substituted or  $L^1$  is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and Wn, where n is 0-3;

wherein each W is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -Q-Ar, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenyl,  $C_3$ - $C_{10}$  cycloalkyl  $C_6$ - $C_{14}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl,

benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by one or more substituents independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup> and halogen up to per-halo; with each R<sup>7</sup> independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenyl,  $C_3$ - $C_{10}$  cycloalkyl  $C_6$ - $C_{14}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by halogen,

wherein Q is -O-, -S-, -N(R<sup>7</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>7</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>- CHX<sup>a</sup>-, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>7</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m= 1-3, and X<sup>a</sup> is halogen; and

Ar is phenyl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -

5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, or 3- or 4-pyridazinyl,pyrazine,

which is optionally substituted by halogen, up to per-halo, and optionally substituted by  $Z_{n1}$ , wherein n1 is 0 to 3 and each Z is independently selected from the group consisting of -CN,  $-CO_2R^7$ ,  $-C(O)R^7$ ,  $-C(O)NR^7R^7$ ,  $-NO_2$ ,  $-OR^7$ ,  $-SR^7$ ,  $-NR^7R^7$ ,  $-NR^7C(O)OR^7$ , -NR<sup>7</sup>C(O)R<sup>7</sup>, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which is C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>2</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkenoyl, C<sub>3</sub>-C<sub>10</sub> cycloalkyl C<sub>6</sub>-C<sub>14</sub> aryl, or 2- or 3-furyl, 2- or 3-thienyl, 2- or 4triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

and optionally substituted by one or more substituents selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and -NR<sup>7</sup>C(O)OR<sup>7</sup>, with R<sup>7</sup> as defined above,

where  $R_g$  is  $C_{1-10}$  alkyl; -CN, -CO<sub>2</sub>R<sub>d</sub>, -OR<sub>d</sub>, -SR<sub>d</sub>, -NO<sub>2</sub>, -C(O)R<sub>e</sub>, -NR<sub>d</sub>R<sub>e</sub>, -NR<sub>d</sub> C(O)OR<sub>e</sub> and -NR<sub>d</sub> C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen,  $C_{1-10}$ , alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl  $C_{6-12}$  aryl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl,

1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

up to per halo substituted  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_3$ - $C_{10}$  cycloalkyl, up to per halo substituted C<sub>6</sub> -C<sub>14</sub> aryl, up to per halo substituted 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl

**Claim 84.** (**Previously presented**) A compound of Formula I:

$$A - D - B \tag{I}$$

or a pharmaceutically acceptable salt thereof, wherein

D is 
$$-NH-C(O)-NH-$$
,

A is a substituted moiety of up to 40 carbon atoms of the formula:  $-L-(M-L^1)_q$ , where L is a 6 membered cyclic structure, which is substituted phenyl or unsubstituted phenyl, bound directly to D,  $L^1$  comprises a substituted cyclic moiety having at least 5 members which is phenyl or pyridinyl,

M is one or more bridging groups selected from the group consisting of -O-, -S-, and -NR<sup>7</sup>- wherein R<sup>7</sup> is hydrogen, q is an integer of from 1-3; and

B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms with at least one 6-member cyclic structure bound directly to D which is phenyl or pyridinyl,

wherein L<sup>1</sup> is substituted by -C(NR<sub>v</sub>)R<sub>z</sub>,

 $R_z$  is hydrogen or a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenyl,  $C_{6-12}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5-, or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

S, N and O, and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-

thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl, C<sub>3</sub>-C<sub>12</sub> halo substituted cycloalkyl up to per halo cycloalkyl halo substituted, up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

or  $-C(O)R_g$ ;

 $R_x$  is  $R_z$  or  $NR_aR_b$  where  $R_a$  and  $R_b$  are

a) independently hydrogen,

a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_1$  - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{2-10}$  alkenyl,  $C_{1-10}$  alkenoyl,  $C_{6-12}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5-, or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{3-12}$  cycloalkyl and is optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-

quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl, C<sub>3</sub>-C<sub>12</sub> halo substituted cycloalkyl up to per halo cycloalkyl, halo substituted, up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

or -C(O)R<sub>g</sub> and are optionally substituted by halogen, or

-OSi( $R_f$ )<sub>3</sub> where  $R_f$  is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O, which is  $C_{1-10}$  alkyl,  $C_{1-10}$  alkoxy,  $C_3$ - $C_{10}$  cycloalkyl  $C_{6-12}$  aryl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 3-, 4-, 5-, 6- or 7-benzotyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 3-

, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl,  $C_3$ - $C_{12}$  halo substituted cycloalkyl up to per halo cycloalkyl, halo substituted, up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzimidazolyl, 1-,

benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or  $-C(O)R_g$ , or

b)  $R_a$  and  $R_b$  together form 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-inidazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

optionally substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-indolyl,

1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_7$  - $C_{24}$  alkaryl,  $C_7$  - $C_{24}$  aralkyl, halo substituted  $C_{1-6}$  alkyl up to per halo alkyl, halo substituted C<sub>6</sub>-C<sub>12</sub> aryl up to per halo aryl, halo substituted C<sub>3</sub>-C<sub>12</sub> cycloalkyl up to per halo cycloalkyl, halo substituted up to per halo, 2- or 3-furyl, 2- or 3thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

or  $-C(O)R_g$ , or

one of  $R_a$  or  $R_b$  is -C(O)-, a  $C_1$ - $C_5$  divalent alkylene group or a substituted  $C_1$ - $C_5$  divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted  $C_1$ - $C_5$  divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, which are  $C_{1-10}$  alkyl,  $C_{3-12}$  cycloalkyl

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-oxazolyl, 2-, 4- or 5-isoxazolyl, 2-

thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

 $C_{1-10}$  alkoxy,  $C_{6-12}$  aryl,  $C_{1-6}$  halo substituted alkyl up to per halo alkyl,  $C_6$ - $C_{12}$  halo substituted aryl up to per halo aryl, C<sub>3</sub>-C<sub>12</sub> halo substituted cycloalkyl, up to per halo cycloalkyl, halo substituted, up to per halo, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5thiazolyl,

or  $-C(O)R_g$ ,

where B is substituted, L is substituted or  $L^1$  is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and Wn, where n is 0-3;

wherein each W is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -C(O)-R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -Q-Ar, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenyl,  $C_3$ - $C_{10}$  cycloalkyl  $C_6$ - $C_{14}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzomidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by one or more substituents independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>C(O)R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup> and halogen up to per-halo; with each R<sup>7</sup> independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which are  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenyl,  $C_3$ - $C_{10}$  cycloalkyl,  $C_6$ - $C_{14}$  aryl,

2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl,

1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by halogen,

wherein Q is -O-, -S-, -N(R<sup>7</sup>)-, -(CH<sub>2</sub>)<sub>m</sub>-, -C(O)-, -CH(OH)-, -(CH<sub>2</sub>)<sub>m</sub>O-, -(CH<sub>2</sub>)<sub>m</sub>S-, -(CH<sub>2</sub>)<sub>m</sub>N(R<sup>7</sup>)-, -O(CH<sub>2</sub>)<sub>m</sub>- CHX<sup>a</sup>-, -CX<sup>a</sup><sub>2</sub>-, -S-(CH<sub>2</sub>)<sub>m</sub>- and -N(R<sup>7</sup>)(CH<sub>2</sub>)<sub>m</sub>-, where m= 1-3, and X<sup>a</sup> is halogen; and

Ar is phenyl, 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, or 3- or 4-pyridazinyl,pyrazine,

which is optionally substituted by halogen, up to per-halo, and optionally substituted by  $Z_{n1}$ , wherein n1 is 0 to 3 and each Z is independently selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -C(O)R<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -NO<sub>2</sub>, -OR<sup>7</sup>, - SR<sup>7</sup> -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)OR<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O, which is  $C_1$ - $C_{10}$  alkyl,  $C_1$ - $C_{10}$  alkoxy,  $C_2$ - $C_{10}$  alkenyl,  $C_1$ - $C_{10}$  alkenoyl,  $C_3$ - $C_{10}$  cycloalkyl  $C_6$ - $C_{14}$  aryl, or 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl,

1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzoxazolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzoxazolyl, 2-, 4-, 5-, 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

and optionally substituted by one or more substituents selected from the group consisting of -CN, -CO<sub>2</sub>R<sup>7</sup>, -COR<sup>7</sup>, -C(O)NR<sup>7</sup>R<sup>7</sup>, -OR<sup>7</sup>, -SR<sup>7</sup>, -NO<sub>2</sub>, -NR<sup>7</sup>R<sup>7</sup>, -NR<sup>7</sup>C(O)R<sup>7</sup>, and -NR<sup>7</sup>C(O)OR<sup>7</sup>, with R<sup>7</sup> as defined above,

where  $R_g$  is  $C_{1-10}$  alkyl; -CN,  $-CO_2R_d$ ,  $-OR_d$ ,  $-SR_d$ ,  $-NO_2$ ,  $-C(O)R_e$ ,  $-NR_dR_e$ , -NR<sub>d</sub> C(O)OR<sub>e</sub> and -NR<sub>d</sub> C(O)R<sub>e</sub>, and R<sub>d</sub> and R<sub>e</sub> are independently selected from the group consisting of hydrogen,  $C_{1-10}$ , alkyl,  $C_{1-10}$  alkoxy,  $C_{3-10}$  cycloalkyl,  $C_{6-12}$  aryl, 2- or 3-furyl, 2or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 1-, 2-, 4- or 5benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7benzisothiazolyl, 2-, 4-, 5-, 6- or 7-benz-1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolinyl, 1-, 3-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl,

up to per halo substituted  $C_1$ - $C_{10}$  alkyl, up to per halo substituted  $C_3$ - $C_{10}$  cycloalkyl, up to per halo substituted  $C_6$ - $C_{14}$  aryl, up to per halo substituted 2- or 3-furyl, 2- or 3-thienyl, 2- or 4-triazinyl, 1-, 2- or 3-pyrrolyl, 1-, 2-, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4-

or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or -5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 2-, 3-, 4-, 5- or 6-2H-thiopyranyl, 2-, 3- or 4-4H-thiopyranyl, 3- or 4-pyridazinyl, pyrazinyl, 2-, 3-, 4-, 5-, 6- or 7-benzofuryl, 2-, 3-, 4-, 5-, 6- or 7-benzothienyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5- 6- or 7-benzisoxazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 2-, 4-, 5-, 6-, 7-, 8- isoquinolinyl, 1-, 2-, 3-, 4- or 9-carbazolyl, 1-, 2-, 3-, 4-, 5-, 6-, 7-, 8- or 9-acridinyl, or 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, or 2- or 3-thienyl, 1,3,4-thiadiazolyl, 3-pyrryl, 3-pyrazolyl, 2-thiazolyl or 5-thiazolyl.

## **REMARKS**

The above amendment cancels claim 61 to eliminate an obviousness type double patenting rejection in a related application (Application No. 09/889,227) and is not responsive to the outstanding office action.

Respectfully submitted,

/Richard J. Traverso/

Richard J. Traverso, Reg. No. 30,595

Attorneys for Applicants

MILLEN, WHITE, ZELANO &
BRANIGAN, P.C.
Arlington Courthouse Plaza 1, Suite 1400
2200 Clarendon Boulevard
Arlington, Virginia 22201

Telephone: (703) 243-6333 Facsimile: (703) 243-6410

Attorney Docket No.: BAYER-15 C1

Date: \_\_\_\_\_ July 23, 2007